

Resonance in an open quantum dot system with a Coulomb interaction: a Bethe-ansatz approach

Akinori NISHINO* and Naomichi HATANO †

Institute of Industrial Science, The University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo, 153-8505

An open quantum system consisting of a quantum dot with a Coulomb interaction and two leads without interactions is studied. The many-body scattering states are constructed with the Bethe-ansatz approach. The expectation value of the electric current is exactly calculated for the scattering states to observe resonance peaks due to many-body scattering.

KEYWORDS: scattering theory, open quantum systems, resonance, interacting resonant-level model, quantum dots, Bethe ansatz

The purpose of this letter is to observe resonance in an open quantum system with a Coulomb interaction. The system that we study is the two-lead interacting resonant-level model (IRLM), which consists of two leads of non-interacting electrons that interact with an electron on a quantum dot in between the two leads. We obtain N -electron scattering states for arbitrary N , generalizing the Bethe-ansatz approach to open systems. By using the scattering states, we exactly calculate the quantum-mechanical expectation value of the electric current through the quantum dot, thereby observing resonance peaks. Some of the resonance peaks appear only when the interaction exists; they reflect the effect of many-body scattering.

The resonance of many-body scattering that we observe in the quantum-mechanical expectation value has not been found in previous works with the Bethe ansatz. The Bethe-ansatz approach has provided a nonperturbative method of studying equilibrium states of interacting quantum systems including the Kondo problem.¹⁻⁵ The approach is now used to discuss transport properties of mesoscopic systems. Konik *et al.*^{6,7} studied transport properties of the Anderson model in the thermodynamic limit of a closed system with *periodic* boundary conditions. Our scattering states, in contrast, appear only in open systems; they are constructed *without* imposing periodic boundary conditions. By extending the Bethe-ansatz approach, Mehta and Andrei⁸ studied the two-lead IRLM as an open system to obtain N -electron scattering states giving nonequilibrium steady states in the limit $N \rightarrow \infty$. In their study, however, the quantum-mechanical expectation value of the current does not depend on the interaction; the effect of the interaction appears only in the statistical-mechanical expectation value as modification of the Fermi distribution in the leads. Thus our results are different from the previous ones.

*E-mail address: nishino@iis.u-tokyo.ac.jp

†E-mail address: hatano@iis.u-tokyo.ac.jp

There has recently been a great deal of interest in mesoscopic systems with interacting electrons. Experiments suggest that interactions are essential in understanding their transport properties.^{9–13} The perturbation theory tells us that the effect of interactions is observed as resonance peaks of the electrical conductance.^{14,15} For non-interacting open quantum systems, the relation between quantum mechanical scattering states and nonequilibrium steady states is well investigated.^{16,17} However, the relation in interacting open quantum systems has not been clarified, excepting Schiller and Hershfield's result¹⁸ at a special point of the interaction parameter where an interacting system is mapped to a non-interacting one. The present study gives a steady step toward an exact analysis of interacting open quantum systems out of equilibrium.

The Hamiltonian of the two-lead IRLM is given by

$$H = \sum_{i=1,2} \left(\int_{-\frac{L}{2}}^{\frac{L}{2}} dz c_i^\dagger(z) \frac{1}{i} \frac{d}{dz} c_i(z) + \frac{t}{\sqrt{2}} (c_i^\dagger(0)d + d^\dagger c_i(0)) \right) + \epsilon_d d^\dagger d + U \sum_{i=1,2} c_i^\dagger(0) c_i(0) d^\dagger d, \quad (1)$$

where $t(>0)$ is the transfer integral between each lead and the quantum dot, ϵ_d is the gate energy of the dot and $U(>0)$ expresses the Coulomb repulsion. The dispersion relation in the leads is linearized in the vicinity of the Fermi energy to be $E = k$, under the assumption that t , ϵ_d and U are small compared with the Fermi energy.^{1,3,5} We stress that we treat the system as an open system in the limit $L \rightarrow \infty$. The one-lead IRLM with periodic boundary conditions was studied with the Bethe ansatz.³ Our purpose is to investigate, for scattering states, the electric current through the quantum dot,

$$I := \dot{N}_2 - \dot{N}_1 = \frac{t}{\sqrt{2}i} \sum_{i=1,2} (-)^i (c_i^\dagger(0)d - d^\dagger c_i(0)). \quad (2)$$

We derive the Schrödinger equations for the system. After the transformation $c_{1/2}(z) = (c_e(z) \pm c_o(z))/\sqrt{2}$, the Hamiltonian (1) is decomposed into the even and odd parts.⁸ Due to the relations $[H, N_e + N_d] = [H, N_o] = 0$ for the number operators $N_{e/o} = \int dz c_{e/o}^\dagger(z) c_{e/o}(z)$ and $N_d = d^\dagger d$, the set $(N_e + N_d, N_o)$ gives a good quantum number. The N -electron state in the subspace with $N_o = n$, ($0 \leq n \leq N$) is generally expressed in the form

$$|\psi\rangle = \left(\int dz g^{(n)}(z) c_e^\dagger(z_1) \cdots c_e^\dagger(z_{N-n}) c_o^\dagger(z_{N-n+1}) \cdots c_o^\dagger(z_N) + \int dz e^{(n)}(z) c_e^\dagger(z_1) \cdots c_e^\dagger(z_{N-n-1}) d^\dagger c_o^\dagger(z_{N-n}) \cdots c_o^\dagger(z_{N-1}) \right) |0\rangle, \quad (3)$$

where $g^{(n)}(z) = g^{(n)}(z_1, \dots, z_N)$ for $0 \leq n \leq N$ and $e^{(n)}(z) = e^{(n)}(z_1, \dots, z_{N-1})$ for $0 \leq n \leq N-1$ are functions to be determined. We also set $e^{(N)}(z) = 0$ for convenience. The eigenvalue

problem $H|\psi\rangle = E|\psi\rangle$ is cast into a set of the Schrödinger equations

$$\begin{aligned}
(N-n) \left(\sum_{i=1}^N \frac{1}{i} \frac{\partial}{\partial z_i} - E \right) g^{(n)}(z) + t \sum_{i=1}^{N-n} (-)^{N-n-i} \delta(z_i) e^{(n)}(\dots, z_{i-1}, z_{i+1}, \dots) &= 0, \\
\left(\sum_{i=1}^N \frac{1}{i} \frac{\partial}{\partial z_i} - E \right) g^{(N)}(z) &= 0, \\
\left(\sum_{i=1}^{N-1} \left(\frac{1}{i} \frac{\partial}{\partial z_i} + U \delta(z_i) \right) + \epsilon_d - E \right) e^{(n)}(z) + t(N-n) g^{(n)}(\dots, z_{N-n-1}, 0, z_{N-n}, \dots) &= 0, \quad (4)
\end{aligned}$$

where $0 \leq n \leq N-1$. In what follows, we use the variables x_i and y_i to express the coordinates of the leads e and o , respectively: $g^{(n)}(z) = g^{(n)}(x_1, \dots, x_{N-n}, y_1, \dots, y_n)$ and $e^{(n)}(z) = e^{(n)}(x_1, \dots, x_{N-n-1}, y_1, \dots, y_n)$. The set of eigenfunctions in the one-electron sector with $E = k$ is given by

$$\begin{aligned}
g^{(0)}(x_1) &= g_k(x_1) := \frac{2e^{ikx_1}}{1 + e^{i\delta_k}} (\theta(-x_1) + e^{i\delta_k} \theta(x_1)), \\
e^{(0)} &= e_k := \frac{t}{k - \epsilon_d}, \\
g^{(1)}(y_1) &= h_k(y_1) := \frac{2e^{iky_1}}{1 + e^{i\delta_k}}
\end{aligned}$$

with the phase shift $\delta_k := -2 \arctan(t^2/2(k - \epsilon_d))$ of one-body scattering at $x_1 = 0$ in the lead e and the step function $\theta(z)$. Note that the eigenfunction $g_k(x_1)$ is discontinuous at $x_1 = 0$.

We construct an N -electron eigenstate with the Bethe ansatz. It is different from the one obtained by Mehta and Andrei.⁸ To demonstrate the difference, we first consider the case $N = 2$. The set of two-electron eigenfunctions with the energy eigenvalue $E = k_1 + k_2$ is assumed to be

$$\begin{aligned}
2g_{k_1 k_2}^{(0)}(x_1, x_2) &= Z_{k_1 k_2}(x_1 - x_2) g_{k_1}(x_1) g_{k_2}(x_2) - Z_{k_1 k_2}(x_2 - x_1) g_{k_2}(x_1) g_{k_1}(x_2), \\
e_{k_1 k_2}^{(0)}(x_1) &= Z_{k_1 k_2}(x_1) g_{k_1}(x_1) e_{k_2} - Z_{k_1 k_2}(-x_1) g_{k_2}(x_1) e_{k_1}, \\
g_{k_1 k_2}^{(1)}(x_1, y_1) &= X(x_1 - y_1) g_{k_1}(x_1) h_{k_2}(y_1), \\
e_{k_1 k_2}^{(1)}(y_1) &= X(-y_1) e_{k_1} h_{k_2}(y_1), \\
2g_{k_1 k_2}^{(2)}(y_1, y_2) &= h_{k_1}(y_1) h_{k_2}(y_2) - h_{k_2}(y_1) h_{k_1}(y_2), \quad (5)
\end{aligned}$$

where the amplitudes $Z_{k_1 k_2}(z)$ and $X(z)$ are defined by

$$\begin{aligned}
Z_{k_1 k_2}(z) &:= e^{-\frac{i}{2} \varphi_{k_1 k_2}} \theta(-z) + e^{\frac{i}{2} \varphi_{k_1 k_2}} \theta(z), \\
X(z) &:= e^{\frac{i}{2} \eta} \theta(-z) + e^{-\frac{i}{2} \eta} \theta(z)
\end{aligned}$$

with the phase shifts $\varphi_{k_1 k_2} := 2 \arctan(-\frac{U}{2} \frac{k_1 - k_2}{k_1 + k_2 - 2\epsilon_d})$ and $\eta := 2 \arctan(-U/2)$ of two-body scattering. The eigenfunctions $g_{k_1 k_2}^{(0)}(x_1, x_2)$ and $e_{k_1 k_2}^{(0)}(x_1)$ are the same Bethe eigenfunctions as those assumed in the one-lead IRLM,³ although we do not impose periodic boundary

conditions. The eigenfunctions $g_{k_1 k_2}^{(1)}(x_1, y_1)$ and $e_{k_1 k_2}^{(1)}(y_1)$ are obtained with separation of variables; if we set $g_{k_1 k_2}^{(1)}(x_1, y_1) = X(x_1 - y_1)\tilde{g}_{k_1 k_2}^{(1)}(x_1, y_1)$ and $e_{k_1 k_2}^{(1)}(y_1) = X(-y_1)\tilde{e}_{k_1 k_2}^{(1)}(y_1)$, Eqs. (4) are decoupled into even and odd parts, and the eigenfunctions $\tilde{g}_{k_1 k_2}^{(1)}(x_1, y_1)$ and $\tilde{e}_{k_1 k_2}^{(1)}(y_1)$ are given by the product of eigenfunctions of the even and the odd parts. The eigenfunctions $g_{k_1 k_2}^{(2)}(y_1, y_2)$ should be a free fermion eigenfunction because of Eq. (4) for $g^{(N)}(z)$. The phase shifts in our solution are different for each two-body scattering in the lead e , in the lead o and between the leads; this gives the resonance of many-body scattering in the current expectation value, as we shall see below. In Mehta and Andrei's solution,⁸ on the other hand, the same phase shift $\varphi_{k_1 k_2}$ of two-body scattering was adopted for all the two-electron eigenfunctions $g^{(n)}(z)$ and $e^{(n)}(z)$.

By exchanging k_1 and k_2 in both eigenfunctions $g_{k_1 k_2}^{(1)}(x_1, y_1)$ and $e_{k_1 k_2}^{(1)}(y_1)$, we have another set of eigenfunctions $g_{k_2 k_1}^{(1)}(x_1, y_1)$ and $e_{k_2 k_1}^{(1)}(y_1)$ with the same eigenvalue $E = k_1 + k_2$. In the limit $t, U \rightarrow 0$, the set $\{g_{k_1 k_2}^{(0)}(x_1, x_2), g_{k_1 k_2}^{(1)}(x_1, y_1), g_{k_2 k_1}^{(1)}(x_1, y_1), g_{k_1 k_2}^{(2)}(y_1, y_2)\}$ reproduces a complete orthogonal system of two free fermions in the two leads, while Mehta and Andrei's solution⁸ does not. In this sense, our solution (5) is more plausible than theirs.

In a way similar to the case $N = 2$, we obtain a set of N -electron eigenfunctions with the energy eigenvalue $E = \sum_{i=1}^N k_i$ in the form

$$g_k^{(n)}(x; y) = \frac{1}{(N-n)!n!} \sum_{\substack{P \in \mathfrak{S}_{N-n} \\ Q \in \mathfrak{S}_n}} \text{sgn}(PQ) Z_{k_P}^{(n)}(x; y) \prod_{i=1}^{N-n} g_{k_{P_i}}(x_i) \prod_{j=1}^n h_{k_{N-n+Q_j}}(y_j),$$

$$e_k^{(n)}(x; y) = \frac{1}{(N-n-1)!n!} \sum_{\substack{P \in \mathfrak{S}_{N-n} \\ Q \in \mathfrak{S}_n}} \text{sgn}(PQ) \tilde{Z}_{k_P}^{(n)}(x; y) \prod_{i=1}^{N-n-1} g_{k_{P_i}}(x_i) e_{k_{P_{N-n}}} \prod_{j=1}^n h_{k_{N-n+Q_j}}(y_j), \quad (6)$$

where $\mathfrak{S}_m$ is the symmetric group acting on the set $\{1, 2, \dots, m\}$ and

$$Z_{k_P}^{(n)}(x; y) = \prod_{1 \leq i < j \leq N-n} Z_{k_{P_i} k_{P_j}}(x_i - x_j) \prod_{\substack{1 \leq i \leq N-n \\ 1 \leq j \leq n}} X(x_i - y_j),$$

$$\tilde{Z}_{k_P}^{(n)}(x; y) = Z_{k_P}^{(n)}(x; y)|_{x_{N-n}=0}.$$

Note that the amplitude $Z_{k_P}^{(n)}(x; y)$ and $\tilde{Z}_{k_P}^{(n)}(x; y)$ are given by the product of the amplitudes of the two-electron eigenfunctions. The N -electron eigenfunctions (6) are indexed by a set of momenta $k = \{k_1, \dots, k_N\}$. We denote by $|k; n\rangle$ the eigenstate obtained by putting the eigenfunctions $g_k^{(n)}(x; y)$ and $e_k^{(n)}(x; y)$ into (3), and call it a Bethe eigenstate.

We show that, for a fixed set of N momenta $k = \{k_1, \dots, k_N\}$, there exist 2^N degenerate Bethe eigenstates with the energy eigenvalue $E = \sum_{i=1}^N k_i$. For a fixed n , we consider ${}_N C_n$ ways of dividing the set k into two subsets wherein the first subset contains $N - n$ elements and the second subset contains n elements. It is convenient to index each way of dividing by $k_R = \{k_{R_1}, \dots, k_{R_N}\}$ with an element R of the symmetric group $\mathfrak{S}_N$ satisfying $R_1 < R_2 < \dots < R_{N-n}$ and $R_{N-n+1} < R_{N-n+2} < \dots < R_N$. The element R is an element of

$\mathfrak{S}_N/(\mathfrak{S}_{N-n} \times \mathfrak{S}_n)$, where $\mathfrak{S}_{N-n}$ is the symmetric group acting on $\{1, 2, \dots, N-n\}$ and $\mathfrak{S}_n$ that acting on $\{N-n+1, N-n+2, \dots, N\}$. For $0 \leq n \leq N$ and $R \in \mathfrak{S}_N/(\mathfrak{S}_{N-n} \times \mathfrak{S}_n)$, all the Bethe eigenstates $|k_R; n\rangle$ with a set of momenta k_R have the same energy eigenvalue $E = \sum_{i=1}^N k_i$. In the limit $L \rightarrow \infty$, the Bethe eigenstates $|k_R; n\rangle$ satisfy the relation

$$\langle k_R; n | k_S; m \rangle = \delta_{nm} \delta_{RS} L^N \prod_{i=1}^N \frac{2}{1 + \cos \delta_{k_i}} + O(L^{N-1})$$

for generic values of $\{k_i\}$. Hence the normalized Bethe eigenstates are orthogonal in the limit $L \rightarrow \infty$. As a result, the total degree of degeneracy of the energy eigenvalue $E = \sum_{i=1}^N k_i$ is $\sum_{n=0}^N {}^N C_n = 2^N$.

We obtain a general N -electron eigenstate by taking a linear combination of the 2^N degenerate Bethe eigenstates $|k_R; n\rangle$ in the form

$$|k\rangle = \sum_{n=0}^N \sum_R \text{sgn}(R) A_R^{(n)} |k_R; n\rangle, \quad (7)$$

where the sum on R runs over elements in $\mathfrak{S}_N/(\mathfrak{S}_{N-n} \times \mathfrak{S}_n)$. The square norm of the eigenstate $|k\rangle$ is readily calculated from $\langle k | k \rangle = \sum_{n=0}^N \sum_R |A_R^{(n)}|^2 \langle k_R; n | k_R; n \rangle$.

The expectation value $\langle I \rangle = \langle k | I | k \rangle / \langle k | k \rangle$ of the current operator I in (2) for each eigenstate $|k\rangle$ in (7) is exactly given by

$$\begin{aligned} \langle I \rangle &= \frac{-t}{2^{N-1} L} \sum_{n=1}^N \frac{n}{(N-n)! n!} \sum_{P \in \mathfrak{S}_N} \text{Im} \left(A_P^{(n)*} A_P^{(n-1)} e^{\frac{i}{2} \delta_{k_{P_{N-n+1}}}} \right) \\ &\quad \times \left(\prod_{i=1}^{N-n} \cos \frac{\varphi_{k_{P_i} k_{P_{N-n+1}}} + \eta}{2} \right) \left(\cos \frac{\eta}{2} \right)^{n-1} \cos \frac{\delta_{k_{P_{N-n+1}}}}{2} e_{k_{P_{N-n+1}}} + O(L^{-2}). \end{aligned} \quad (8)$$

Here, by using the fact that any element $P \in \mathfrak{S}_N$ is decomposed as $P = RQ$ with a unique element $R \in \mathfrak{S}_N/(\mathfrak{S}_{N-n} \times \mathfrak{S}_n)$ and a unique element $Q \in \mathfrak{S}_{N-n} \times \mathfrak{S}_n$, we set $A_P^{(n)} = A_R^{(n)}$ for every P with the same R .

The result in (8) shows that we need to consider the linear combination (7) of the degenerate Bethe states in order to obtain a non-zero current expectation value. Indeed, a specific Bethe eigenstate $|k_R; n\rangle$ gives the expectation value $\langle k_R; n | I | k_R; n \rangle = 0$. We stress that we do not impose periodic boundary conditions to the eigenfunctions $g_k^{(n)}(x; y)$ and $e_k^{(n)}(x; y)$ in (6); with the periodic boundary conditions,^{6,7} the eigenstates $|k_R; n\rangle$ with different n or different R would not be degenerate, and hence the current expectation value would be zero.

By expressing the eigenstate $|k\rangle$ in terms of the leads 1 and 2, the eigenfunction describing $N-n$ electrons in the lead 1 and n electrons in the lead 2 is given by

$$\begin{aligned} (N-n)! n! F_k^{(n)}(z) &:= \langle c_2(z_N) \cdots c_2(z_{N-n+1}) c_1(z_{N-n}) \cdots c_1(z_1) | k \rangle \\ &= \frac{1}{2^{\frac{N}{2}}} \sum_{m=0}^N \sum_R (-)^{\sharp\{R_i | N-n < R_{N-m+1}, \dots, R_N\}} \text{sgn}(R) \end{aligned}$$

$$\times \langle c_o(z_{R_N}) \cdots c_o(z_{R_{N-m+1}}) c_e(z_{R_{N-m}}) \cdots c_e(z_{R_1}) | k \rangle,$$

where $\#A$ stands for the number of elements in the set A . We consider the behavior of $F_k^{(n)}(z)$ in the region $z_1 < z_2 < \cdots < z_N < 0$. The eigenfunction $F_k^{(n)}(z)$ is a complicated linear combination of plane waves $e^{i \sum_i k_{P_i} z_i}$ for $P \in \mathfrak{S}_N$. Among them, we call the plain wave $e^{i \sum_i k_i z_i}$ an “incoming wave”. The terms with the incoming wave are summarized as

$$e^{i \sum_i k_i z_i} \sum_{m,R} (-)^{\# \{R_i | N-n < R_{N-m+1}, \dots, R_N\}} \tilde{A}_R^{(m)}$$

where

$$\tilde{A}_R^{(m)} = A_R^{(m)} e^{-\frac{i}{2} (\sum_{i < j \leq N-m} \varphi_{k_{R_i} k_{R_j}} + \eta \sum_{\substack{i \leq N-m \\ N-m < j}} \text{sgn}(R_i - R_j))}.$$

We define the scattering states $|k\rangle^{(\ell)}$, $(\ell = 0, 1, \dots, N)$ by taking the coefficients $\{\tilde{A}_R^{(n)}\}$ of the eigenstate $|k\rangle$ as

$$\tilde{A}_R^{(n)} = (-)^{\# \{R_i | N-\ell < R_{N-n+1}, \dots, R_N\}}, \quad (9)$$

for $k_1 > k_2 > \cdots > k_N$. In the scattering state $|k\rangle^{(\ell)}$, the incoming wave $e^{i \sum_{i=1}^N k_i z_i}$ exists only in the eigenfunction $F_k^{(\ell)}(z)$, which describes $N - \ell$ electrons in the lead 1 and ℓ electrons in the lead 2.

The scattering states $|k\rangle^{(\ell)}$ are different from those of the standard one-body scattering theory in quantum mechanics. If we were solving the one-body scattering problem, the scattering state would be obtained from the condition that an electron comes only from the lead 1 or the lead 2. However, the eigenstate $|k\rangle$ in (7) does not give such scattering state. In fact, the scattering state $|k\rangle^{(\ell)}$ extends to all parts of the two leads for $U > 0$. In other words, it is impossible to judge whether each electron comes from the lead 1 or the lead 2 for $U > 0$, which is not strange since we assume the same Fermi energy for both leads. In the limit $U \rightarrow 0$, our scattering state $|k\rangle^{(\ell)}$ is reduced to the standard one-body scattering state.

By applying (9) to the expectation value (8), we have

$$\begin{aligned} \langle I \rangle^{(\ell)} &= \frac{t}{2^{N-1} L} \sum_{n=1}^N \frac{n}{(N-n)! n!} \sum_{P \in \mathfrak{S}_N} \text{sgn}(P_n - N + \ell) \\ &\times \sin \left(\frac{1}{2} \left(\sum_{i=1}^{n-1} \text{sgn}(P_i - P_n) \varphi_{k_{P_i} k_{P_n}} + \sum_{i=1}^N \text{sgn}(P_i - P_n) \eta - \delta_{k_{P_n}} \right) \right) \\ &\times \left(\prod_{i=1}^{n-1} \cos \frac{\varphi_{k_{P_i} k_{P_n}} + \eta}{2} \right) \left(\cos \frac{\eta}{2} \right)^{N-n} \cos \frac{\delta_{k_{P_n}}}{2} e_{k_{P_n}} + O(L^{-2}). \end{aligned}$$

Short calculations reveal that every term in $\langle I \rangle^{(\ell)}$ contains the product of the factors $\sin \delta_k$, $1 + \cos \delta_k$, $\sin(\varphi_{k_i k_j} + \eta)$ and $1 + \cos(\varphi_{k_i k_j} + \eta)$, which are rational functions of k_i , t , ϵ_d and U . The factors have poles at $\epsilon_d = k_i \pm it^2/2$, $(k_i + k_j)/2 \mp i(k_i - k_j)U/4$ in the complex plane of ϵ_d . Figure 1 shows the current expectation value $\langle I \rangle^{(\ell)}$ as a function of the gate energy ϵ_d for the scattering states $|k\rangle^{(\ell)}$ indexed by $(N, \ell) = (2, 0)$ and $(3, 1)$.




Fig. 1. The current expectation value $\langle I \rangle^{(\ell)}$ for the scattering states. We fixed $t = 1$.

We find resonance peaks in the vicinity of $\epsilon_d = (k_i + k_j)/2$, which correspond to many-body scattering; they appear only for $U > 0$. As is stressed above, the resonance of many-body scattering is originated from the phase shifts which are different for each two-body scattering in the lead e , in the lead o and between the two leads. We also find resonance peaks in the vicinity of $\epsilon_d = k_i$, which correspond to one-body scattering at the quantum dot and are reduced to Lorentzian peaks in the limit $U \rightarrow 0$. The resonance peaks in the vicinity of $\epsilon_d = (k_i + k_j)/2$ were not present in Mehta and Andrei's result;⁸ their results are equal to the limit $U \rightarrow 0$ of our result. This is because the interaction effect would be canceled in the current expectation value $\langle I \rangle$ if we adopted the same phase shifts for all the two-body scattering in the lead e , in the lead o and between the two leads. Our choice (5) of the phase shifts of two-body scattering is more plausible in the context of eigenstates as mentioned above. It would be interesting to discuss how the resonance of many-body scattering affects the transport properties of the interacting open quantum system out of equilibrium.

Acknowledgments

The authors would like to thank Prof. T. Deguchi, Dr. T. Imamura, Dr. K. Sasada and Dr. M. Matsuo for helpful comments. The present study is partially supported by Core Research for Evolutional Science and Technology of Japan Science and Technology Agency.

References

- 1) N. Andrei, Phys. Rev. Lett. **45** (1980), 379.
- 2) N. Andrei, K. Furuya and J. H. Lowenstein, Rev. Mod. Phys. **55** (1983), 331.
- 3) V. M. Filyov and P. B. Wiegmann, Phys. Lett. A **76** (1980), 283.
- 4) A. M. Tsvelick and P. B. Wiegmann, Adv. Phys. **32** (1983), 453.
- 5) P. B. Wiegmann, Phys. Let. A **80** (1980), 163.
- 6) R. M. Konik, H. Saleur and A. Ludwig, Phys. Rev. B **66** (2002), 125304.
- 7) R. M. Konik, H. Saleur and A. W. W. Ludwig, Phys. Rev. Lett. **87** (2001), 236801.
- 8) P. Mehta and N. Andrei, Phys. Rev. Lett. **96** (2006), 216802.
- 9) S. M. Cronenwett, T. H. Oosterkamp and L. P. Kouwenhoven, Science **281** (1998), 540.
- 10) D. Goldhaber-Gordon, J. Gores, M. A. Kastner, H. Shtrikman, D. Mahalu and U. Meirav, Phys. Rev. Lett. **81** (1998), 5225.
- 11) D. Goldhaber-Gordon, H. Shtrikman, D. Mahalu, D. Abusch-Magder, U. Meirav and M. A. Kastner, Nature (London) **391** (1998), 156.
- 12) D. C. Ralph and R. A. Buhrman, Phys. Rev. Lett. **69** (1992), 2118.
- 13) D. C. Ralph and R. A. Buhrman, Phys. Rev. Lett. **72** (1994), 3401.
- 14) Y. Meir, N. S. Wingreen and P. A. Lee, Phys. Rev. Lett. **66** (1991), 3048.
- 15) A. Levy Yeyati, A. Martin-Rodero and F. Flores, Phys. Rev. Lett. **71** (1993), 2991.
- 16) S. Datta, Electronic Transport in Mesoscopic Systems (Cambridge University, Cambridge, England, 1995).
- 17) A. M. Zagoskin, Quantum Theory of Many-Body Systems, Graduate Texts in Contemporary Physics (Springer, New York, 1998).
- 18) A. Schiller and S. Hershfield, Phys. Rev. B **58** (1998), 14978.